

WARNING

KEEP OUT OF REACH OF CHILDREN



Advion[®] Cockroach Gel

Insecticide

syngenta.

Active Ingredient:

ADVION COCKROACH GEL contains 6 g/kg INDOXACARB in the form of a gel.

For the control of Cockroaches in the situations specified in the Directions for Use

Net Contents: 30 g

Approved pursuant to the HSNO Act 1996, Approval Code HSR001752

Syngenta Crop Protection Limited

Tower 2, Level 7, 110 Symonds Street, Auckland

In a transport emergency dial 111, Police or Fire Brigade.

For specialist advice in an emergency only, call 0800 734 607 (24 hours).

TM

PLEASE READ THE LABEL COMPLETELY BEFORE USE

This product **MUST NOT** be used for any purpose or in any manner contrary to the controls and conditions of this registration.

HSNO Classifications 6.9B, 9.1C, 9.4B

**WARNING**

May cause organ damage (hematopoietic system) from repeated oral exposure at high doses.
Harmful to aquatic organisms with long lasting effects.
Toxic to terrestrial invertebrates.

PRECAUTIONS

Store in the closed, original container in a cool, well-ventilated locked place out of the reach of children.
DO NOT store in direct sunlight and DO NOT store at temperatures exceeding 35°C.
Avoid contact with skin and eyes.
If clothing becomes contaminated with product, remove clothing immediately.
DO NOT eat, drink or smoke while using.
Wash hands and face before eating, drinking or smoking and after work.
DO NOT apply to surfaces that food contacts.
DO NOT allow open foods to contact gel material.
Avoid contamination of any water supply with chemical or empty container.

Container disposal: Ensure the container is empty. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in the garbage. DO NOT reuse this container for any other purpose.

Product disposal: Dispose of this product only by using according to the label, or as above.

FIRST AID

If swallowed DO NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. Begin artificial respiration if breathing has stopped. Use mouth to nose rather than mouth to mouth. Obtain medical attention. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin. If splashed in the eyes, wash out immediately with water. If inhaled, move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

SPILLS

Wear suitable protective clothing including face shield, impervious gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

GENERAL INFORMATION

ADVION COCKROACH GEL is designed to control pest species of cockroaches including German and American. The slight delay in mortality caused by the active ingredient, Indoxacarb, allows cockroaches to consume the bait and return to the harbourage site to contaminate other cockroaches resulting in significant reduction in infestation levels.

DIRECTIONS FOR USE

Restraints:

DO NOT place gel on surfaces that are regularly washed.

DO NOT allow to contact food, livestock feed or their water.

Before using all unpackaged food and packaging material must be removed from the room.

DO NOT place gel on surfaces where food is prepared or on utensils that are used for food preparation, on food packing materials or on surfaces where food packing materials are stored.

ADVION Cockroach Gel may be used in cracks and crevices as a cockroach pesticide in conjunction with an approved vermin control program provided its use does not create a nuisance.

| Situation | Pest | Rates | Critical Comments |
|---|--|---|---|
| Domestic, Commercial and Public Service buildings including; Farm buildings, Food processing establishments, Factories, Hospitals, Homes, Kitchens, Offices, Restaurants, Retail outlets, Store-rooms, Vehicles and other situations which fall within these definitions. | German cockroach (<i>Blattella germanica</i>), American cockroach (<i>Periplaneta americana</i>) and other common cockroaches. | Apply at least 2 spots per m ² . Increasing the number of spots (up to 5/m ²) may be required for heavy infestations or if American cockroaches are the main pest. Recommended spot size is approximately 5 mm diameter (0.1 g). | Infested areas should be carefully inspected to determine overall infestation levels, locations of harbourage areas, and most appropriate application points. ADVION COCKROACH GEL should be applied as a small spot of material, approximately 0.1 grams (approx. 5 mm in diameter - the size of a small pea), to areas not able to be contacted by children or pets. Several small applications distributed within an area are more effective than 1 or 2 applications of large quantities. In some cases a small thin bead (less than 2.5 mm wide and 5 cm long) may be appropriate such as between construction elements or other cracks and crevices. Indoor applications can include cracks and crevices along walls or floors, behind or under equipment, under tables, within framing of tables or furniture, around sinks, within or near garbage collection areas, attics, crawl spaces, and cracks and crevices in cabinets. Bait placements should be inspected periodically for consumption or attractiveness and reapplied as needed. A typical infestation may require as little as 1/2 a tube (15 grams) of ADVION COCKROACH GEL whereas heavy infestations in large structures may require the use of several tubes of gel bait. |

NZFSA Approved Type All animal product except dairy

AgriQuality approved for food, beverage/dairy farm & factory food areas non contact (crevice treatment) ref # 1954

APPLICATION

1. ADVION cockroach gel bait should be applied to areas frequented by cockroaches such as between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, and behind and under appliances, cabinets, sinks, closets, and any other equipment that cockroaches are typically known to hide.
2. Avoid applying Advion cockroach gel bait to surfaces recently treated with residual sprays.
3. Surfaces treated with Advion cockroach gel bait should not be treated with residual sprays.
4. Repeat applications may be made upon inspection during follow-up inspections of the treated areas.

To apply the gel, remove the cap on the nozzle and touch the tip to the surface to be treated and slowly depress plunger until sufficient gel is dispensed. Recap the dispenser after treatment is completed.

Where maintenance treatments form part of the professional pest controllers contract it is recommended that cockroach activity be monitored where gel has been applied one month after initial treatment or at more regular intervals in areas of high infestation.

STORAGE

Store in the closed, original container in a cool, well-ventilated locked place out of the reach of children. DO NOT store in direct sunlight. DO NOT store at temperatures exceeding 35°C. See Safety Data Sheet for additional storage information. Stores containing quantities 1000 kg or more require signage and secondary containment.

SAFETY DATA SHEET

If additional hazard information is required, refer to the Safety Data Sheet. For a copy, visit our website at www.syngenta.co.nz

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company



Registered to and distributed by Syngenta Crop Protection Limited, Tower 2, Level 7, 110 Symonds Street, Auckland.

Bar code

Date of manufacture

Batch number