

The ultimate pre-emergent BARRICADE for turf grass and garden beds




syngenta®

BARRICADE Herbicide controls a wide range of pre-emergent grassy weeds and importantly breaks the germination cycle of *Poa annua*. You can trust BARRICADE to last for up to 6 months so your application window is more flexible.

BARRICADE binds high in the soil profile ensuring turf and ornamental roots below the barrier are not affected and your turf and ornamentals stay in prime condition.

What BARRICADE offers you:

- Up to six months pre-emergent control of grassy weeds in turf including African Lovegrass, Parramatta Grass, Summer Grass, Crab Grass, Crowsfoot Grass and Winter Grass
- Low odour and non-staining formulation
- Flexible application window
- Registered for use in garden beds for broadleaf and grassy weed control
- Easy to apply spray over the top of turf and ornamentals



BARRICADE can be used on the following established turf species:

Warm Season Turf Grasses

Bahia Grass, Buffalo Grass, Carpet Grass, Common Couch, Hybrid Couch, Kikuyu, Queensland Blue Couch, Seashore Paspalum, Zoysia

Refer to the registered label for full details.

SpraySURE | Accuracy and Performance

To achieve great results, accurate application is important.

Target	Syngenta Nozzle	Optimal Water Rate	Optimal Pressure Range
Root and soil So	XC 08	> 500 L/ha	2 to 4 bar

BARRICADE can deliver up to 6 months residual control. Positioning of applications can be carried out well in advance of weeds germinating and still deliver season long control.

Example: Spring applications for the control of summer grassy weeds like *Digitaria* (Summer Grass and Crab Grass) and *Eleusine* (Crowsfoot Grass) should ideally be applied around the end of August. Control will last up to the end of February. Autumn applications for the control of *Poa* (Winter Grass) should ideally be applied around the end of February. Control will last up to the end of August.

BARRICADE as a pre-emergent herbicide works in the top soil layers. Application targeted towards delivering BARRICADE close to the soil surface is best suited to ensure optimum results. BARRICADE needs to be washed in within 7 days after application.

Target Weeds	Rate (Turf)	Rate (Garden Beds)	SpraySure
Barnyard Grass, Crab Grass, Crowsfoot Grass, Rat's Tail, Summer Grass, Vulpia Hair Grass	1 to 3 L/ha	4 to 8 L/ha	So
African Lovegrass, Bahia Grass, Kentucky Blue Grass	2 to 3 L/ha	4 to 8 L/ha	So
Paspalum	2 to 4 L/ha	4 to 8 L/ha	So
Winter Grass	1 to 4 L/ha	4 to 8 L/ha	So
Chickweed, Cudweed, Oxalis, Prairie Grass, Ryegrass, Sow Thistle, Willowherb	n/a	4 to 8 L/ha	So

Refer to the registered label for full details and instructions for use. **F** = Foliar applied **So** = Soil applied **Su** = Surfactant required

Active Ingredient

BARRICADE contains 476.2 g/L prodiamine, a member of the dinitroaniline herbicides.

GROUP **K1** HERBICIDE

Formulation and Packaging

5 L pack size

A suspension concentrate (SC) with minimal odour and non-staining properties. BARRICADE mixes completely with water and produces a homogenous easy to apply suspension.



www.syngenta.co.nz
www.greencast.com.au

 **Barricade**[®]
Herbicide

syngenta[®]

PGG Wrightson Turf

For more information contact PGG Wrightson Turf on (09) 570 2570 (North Island) or (03) 372 8719 (South Island).

Syngenta Crop Protection Limited, Tower 2, Level 7, 110 Symonds St, Auckland. Phone: (09) 306 1500 Fax: (09) 306 1501, www.syngenta.co.nz. [®] Registered trademark of a Syngenta Group Company. [™] Trademark of a Syngenta Group Company. All products written in uppercase are registered trademarks of a Syngenta Group Company. ©Syngenta 2017.

PLEASE READ LABEL COMPLETELY BEFORE USE.

Barricade approved pursuant to the HSNO Act 1996, Approval Code HSR101240. SB 17-347