

New Zealand

Published on Syngenta (<https://www.syngenta.co.nz>)

[Home](#) > Dovetail

---

## Dovetail



# Dovetail

Last updated:

13.11.2018

## Insecticide

**Authorisation Number:**

ACVM: P7941 | HSNO: HSR008052

**Pack size:**

5 litres

**Composition:**

5 g/L

LAMBDA-CYHALOTHRIN

100 g/L

PIRIMICARB

**Formulation:**

Emulsifiable Concentrate

DOVETAIL is an insecticide for insect control in Cereals, Brassicas and Potatoes.

## Product CP: Tabs

- [Registrations](#)
- [Classification and Security](#)
- [Application Advice](#)

## [Registrations](#)

## Dovetail Cereals

Cereals

Ensure sufficient water volume is used to give thorough crop penetration and coverage.

### **Aphids**

<b>Recommended dose</b>	<b>Maximum number of applications</b>	<b>Pre Harvest Interval (PHI)</b>
1 L/ha	2 per season	Grain: 70 Days. Forage: 28 Days
<b>Stage of use:</b>	Apply at first sign of aphids. A second application may be needed if aphids continue to migrate into the crop.	

---

## Dovetail Forage Brassicas

Forage Brassicas

Add a non-ionic surfactant at label rates when treating brassicas.

### **Aphids**

<b>Recommended dose</b>	<b>Maximum number of applications</b>	<b>Pre Harvest Interval (PHI)</b>
1 to 1.25 L in 400 to 600 litres	Refer to label	Refer to label
<b>Stage of use:</b>	Application to Forage Brassicas should be made prior to a build up of pest numbers and before significant damage occurs.	

---

### **Diamondback Moth**

<b>Recommended dose</b>	<b>Maximum number of applications</b>	<b>Pre Harvest Interval (PHI)</b>
1 to 1.25 L in 400 to 600 litres	Refer to label	Refer to label
<b>Stage of use:</b>	Application to Forage Brassicas should be made prior to a build up of pest numbers and before significant damage occurs.	

---

### **White Butterfly**

<b>Recommended dose</b>	<b>Maximum number of applications</b>	<b>Pre Harvest Interval (PHI)</b>
1 to 1.25 L in 400 to 600 litres	Refer to label	Refer to label
<b>Stage of use:</b>	Application to Forage Brassicas should be made prior to a build up of pest numbers and before significant damage occurs.	

---

## Dovetail Potatoes

Potatoes

For Tuber Moth control it is important to maintain adequate soil moisture and soil cover over developing tubers to assist prevention of tuber mining.

### Aphids

<b>Recommended dose</b>	<b>Maximum number of applications</b>	<b>Pre Harvest Interval (PHI)</b>
-------------------------	---------------------------------------	-----------------------------------

1.5 L in 200 to 400 litres	Refer to label	14 Days
----------------------------	----------------	---------

**Stage of use:** Apply at first sighting of insect pests then at two weekly intervals or as indicated by pest activity.

---

### Potato Tuber Moth

<b>Recommended dose</b>	<b>Maximum number of applications</b>	<b>Pre Harvest Interval (PHI)</b>
-------------------------	---------------------------------------	-----------------------------------

1.5 L in 200 to 400 litres	Refer to label	14 Days
----------------------------	----------------	---------

**Stage of use:** Apply at first sighting of insect pests then at two weekly intervals or as indicated by pest activity.

---

## Dovetail Vegetable Brassicas

Vegetable Brassicas

Refer to label

### Aphids

<b>Recommended dose</b>	<b>Maximum number of applications</b>	<b>Pre Harvest Interval (PHI)</b>
-------------------------	---------------------------------------	-----------------------------------

1 to 1.25 L/ha	Refer to label	3 Days
----------------	----------------	--------

**Stage of use:** For Vegetable Brassicas apply to seedbeds and following transplanting at two week intervals or as indicated by pest activity.

---

### Diamondback Moth

<b>Recommended dose</b>	<b>Maximum number of applications</b>	<b>Pre Harvest Interval (PHI)</b>
-------------------------	---------------------------------------	-----------------------------------

1 to 1.25 L in 400 to 600 litres	Refer to label	3 Days
----------------------------------	----------------	--------

**Stage of use:** For Vegetable Brassicas apply to seedbeds and following transplanting at two week intervals or as indicated by pest activity.

---

### White Butterfly

<b>Recommended dose</b>	<b>Maximum number of applications</b>	<b>Pre Harvest Interval (PHI)</b>
1 to 1.25 L in 400 to 600 litres	Refer to label	3 Days
<b>Stage of use:</b>	For Vegetable Brassicas apply to seedbeds and following transplanting at two week intervals or as indicated by pest activity.	

## **Classification and Security**

### **Re-entry Period**

- DO NOT enter treated area within 24 hours without protective clothing until spray has dried.

### **Hazard Statements**

- 3.1D
- 6.1D
- 6.1E
- 8.2C
- 8.3A
- 6.5B
- 6.9B
- 9.1A
- 9.2C
- 9.3B
- 9.4C

### **Personal Protective Equipment**

- When mixing, loading or applying wear protective clothing including chemical resistant coveralls, chemical resistant gloves, chemical resistant footwear plus socks and chemical resistant headgear for overhead exposures.

## **Application Advice**

### **Tank mixing**

Half fill spray tank with clean water, and add the required amount of DOVETAIL (and adjuvant where recommended) and agitate during completion of filling. Maintain agitation while spraying.

### **Water volume**

Apply in sufficient water to ensure thorough and uniform coverage.

### **Rainfastness**

2 hours